

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 745 596 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

04.12.1996 Bulletin 1996/49

(21) Application number: 95940466.6

(22) Date of filing: 18.12.1995

(51) Int. Cl.⁶: C07D 263/32, C07D 413/04,
A61K 31/42, C07D 207/333,
C07D 265/30, C07D 295/10,
C07D 307/46(86) International application number:
PCT/JP95/02600(87) International publication number:
WO 96/19463 (27.06.1996 Gazette 1996/29)

(84) Designated Contracting States:

AT BE CH DE DK ES FR GB GR IE IT LI NL PT SE

(30) Priority: 20.12.1994 JP 335838/94

27.03.1995 JP 93099/95

06.06.1995 JP 164656/95

20.11.1995 JP 326571/95

(71) Applicant: Japan Tobacco Inc.

Minato-Ku Tokyo 105 (JP)

(72) Inventors:

• HARUTA, Junichi



Central Pharm. Research Institute

Takatsuki-shi Osaka 569 (JP)

• HASHIMOTO, Hiromasa
Central Pharm. Research Inst.
Takatsuki-shi Osaka 569 (JP)• MATSUSHITA, Mutsuyoshi
Cent. Pharm. Research Inst.
Takatsuki-shi Osaka 569 (JP)(74) Representative: von Kreisler, Alek, Dipl.-Chem. et
al
Patentanwälte,
von Kreisler-Selting-Werner,
Bahnhofsvorplatz 1 (Deichmannhaus)
50667 Köln (DE)

(54) HETEROAROMATIC OXAZOLE COMPOUNDS AND USE THEREOF

(57) A heterocyclic aromatic oxazole compound of the formula (I)

wherein Z is an oxygen atom; one of R and R₁ is a group of the formulawherein R₃ is lower alkyl, amino or lower alkylamino, and R₄, R₅, R₆ and R₇ are the same or different and each is hydro-

EP 0 745 596 A1

gen atom, halogen atom, lower alkyl, lower alkoxy, trifluoromethyl, hydroxy or amino, provided that at least one of R_4 , R_5 , R_6 and R_7 is not hydrogen atom, and the other is an optionally substituted cycloalkyl, an optionally substituted heterocyclic group or an optionally substituted aryl; and R_2 is a lower alkyl or a halogenated lower alkyl, and a pharmaceutically acceptable salt thereof. The heterocyclic aromatic oxazole compound and pharmaceutically acceptable salts thereof have antipyretic action, analgesic action, anti-inflammatory action, and particularly, selective inhibitory action on cyclooxygenase-2 (COX-2), and are expected to be useful as anti-inflammatory agents with less side-effects such as digestive tract disorders.

Description

Technical Field

The present invention relates to novel heterocyclic aromatic oxazole compounds. More particularly, the present invention relates to heterocyclic aromatic oxazole compounds having antipyretic activity, analgesic activity, anti-inflammatory activity, and in particular, selective inhibitory activity against cyclooxygenase-2 (COX-2), pharmaceutically acceptable salts thereof, intermediates for producing them and pharmaceuticals useful as anti-inflammatory agents causing less side-effects such as disorders in the digestive tract, which comprise these heterocyclic aromatic oxazole compounds.

Background Art

It has been conventionally known that arachidonic acid metabolites, prostaglandin E₂ (PGE₂), prostaglandin I₂ (PGI₂) and thromboxane B₂ (TXB₂) are deeply involved in inflammations. An important enzyme in this arachidonic acid metabolism is cyclooxygenase. Cyclooxygenase is a synthase which produces prostaglandin H₂ (PGH₂) from arachidonic acid via prostaglandin G₂ (PGG₂), and includes cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2).

With respect to COX-1, cDNA cloning was performed in 1988 and its primary structure and induction by various factors have been clarified [Yokoyama, C. et al.: Biochem. Biophys. Res. Commun., 165: 888-894 (1989); Smith, W. L. et al.: Biochim. Biophys. Acta, 1083: 1-17 (1991); DeWitt, D. L.: Biochim. Biophys. Acta, 1083: 121-134 (1991)]. On the other hand, the existence of an isozyme of COX-1, namely, COX-2, was suggested in 1989 [Holtzman, M. J. et al.: J. Biol. Chem., 267: 21438-21445 (1992)], and cDNAs of COX-2 of chicken, mouse and human have been cloned since 1991 [Xie, W. et al.: Proc. Natl. Acad. Sci. USA, 88: 2692-2696 (1991); Kujubu, D. A. et al.: J. Biol. Chem., 266: 12866-12872 (1991); Hla, T. et al.: Proc. Natl. Acad. Sci. USA, 89: 7384-7388 (1992)]. COX-2 is quickly induced by phorbol ester, lipopolysaccharide (LPS) and the like, and the relationship with inflammation and bronchial asthma has been inferred.

COX-1 systemically and constantly exists in almost all cells and is physiologically concerned with the generation of prostaglandin (PG) necessary for the functions of, for example, stomach and kidney. Therefore, when COX-1 is inhibited, the biosynthesis of PG by vasodilative PGE₂ and PGI₂, which protect gastric mucosa, is suppressed, and the protective action on the gastric mucosa becomes degraded, as a result of which ulcer is caused. With regard to a symptom associated with a decrease in renal blood flow, in general terms, the renal blood flow can be increased by promoting the production of vasodilative PGE₂ in the body, thereby to appropriately maintain glomerular filtration rate. However, if the production of such vasodilative PG is suppressed due to the inhibition of COX-1, the renal blood flow becomes less, so that a side-effect such as the onset of ischemic acute renal insufficiency is sometimes caused.

On the other hand, COX-2 exists in particular sites such as monocytes, synovial cells, granulosa cells and intravenous endothelial cells, and is topically expressed when inflammation is caused. It is therefore considered that PG generated by COX-2 is deeply concerned with inflammation and tissue disorders.

Currently, non-steroidal anti-inflammatory drugs (NSAID) such as aspirin, mefenamic acid, diclofenac, indomethacin, ibuprofen and naproxen have been widely used in clinical situations. Most of these NSAIDs are anti-inflammatory drugs which selectively inhibit cyclooxygenase (COX) and are associated with side-effects such as disorders in the digestive tract. Such side-effects are considered to be caused by the fact that they, though certainly selectively inhibit COX, inhibit both COX-1 and COX-2.

It follows therefrom that selective inhibition, without inhibition of COX-1, of solely COX-2 which is specifically induced at the inflammatory sites, would enable provision of a superior anti-inflammatory drug free of side-effects such as disorders in the digestive tract (e.g., ulcer).

There are various reports on anti-inflammatory drugs having selective COX-2 inhibitory activity, which aim at reducing side-effects such as disorders in the digestive tract.

For example, WO94/15932 discloses, as COX-2 inhibitors, 5-membered heterocyclic compounds substituted by bisaryl, such as thiophene, furan and pyrrole, which are specifically exemplified by 3-(4-methylsulfonylphenyl)-4-(4-fluorophenyl)thiophene. However, this publication merely shows a 5-membered heterocyclic compound such as thiophene having aryl or heteroaryl at the 3-position or 4-position.

Moreover, various reports deal with anti-inflammatory drugs having cyclooxygenase-inhibitory action, prostaglandin synthesis-inhibitory action or thromboxane A₂ synthesis-inhibitory action.

For example, Japanese Patent Unexamined Publication No. 141261/1991 discloses pyrazole derivatives such as ethyl 1-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]pyrazole-3-carboxylate; Japanese Patent Unexamined Publication No. 183767/1982 discloses thiazole derivatives such as 2-methylthio-5-phenyl-4-(3-pyridyl)thiazole; and Japanese Patent Unexamined Publication No. 58981/1985 discloses thiazole derivatives such as 2-ethyl-4-(4-methoxyphenyl)-5-(3-pyridyl)-1,3-thiazole. These publications mention that they are useful as anti-inflammatory drugs, whereas they do not disclose if they have selective inhibitory action on COX-2 to reduce side-effects, or any suggestion of it.

There are other reports on the following heterocyclic aromatic compounds.

For example, US Patent No. 4632930 discloses oxazole compounds such as 5-cyclohexyl-4-(4-methylsulfonylphenyl)- α,α -bis(trifluoromethyl)oxazole-2-methanol. Yet, the compounds disclosed therein are effective for hypertension and their usefulness as anti-inflammatory drugs or any suggestion to that effect are not included.

Japanese Patent Application under PCT laid-open under Kohyo No. 500054/1984 discloses oxazole derivatives having heteroaryl or carbon ring aryl at the 4-position or 5-position of oxazole ring and having carboxy, ester or amidized carboxy via lower alkylene at the 2-position thereof, such as ethyl 2-[4-phenyl-5-(3-pyridyl)-oxazol-2-yl]-propionate; and Japanese Patent Application under PCT laid-open under Kohyo No. 500055/1984 discloses imidazole derivatives having heteroaryl and/or carbon ring aryl at the 4-position or 5-position of imidazole ring and having formyl or acetalized formyl via lower alkylene at the 2-position thereof, such as 2-[4-phenyl-5-(3-pyridyl)-imidazol-2-yl]-acetaldehyde dimethyl acetal. These publications teach that these compounds are effective as dermal antiphlogistic or mucosal antiphlogistic for inflammatory dermal diseases, but do not teach or even suggest that they have selective inhibitory action on COX-2.

Japanese Patent Unexamined Publication No. 70446/1993 discloses N-thiazolylsulfonamide derivatives such as N-[5-cyclohexyl-4-(4-methoxyphenyl)thiazol-2-yl]trifluoromethanesulfonamide; and Japanese Patent Unexamined Publication No. 83372/1990 discloses cyclohexylimidazole derivatives such as 4-cyclohexyl-5-phenyl-2-t-butyl-imidazole. These publications only exemplify cyclohexyl as a substituent and include no suggestion as to the substitution with phenyl substituted by aminosulfonyl, lower alkylaminosulfonyl or lower alkylsulfonyl.

WO94/27980 discloses oxazole compounds such as 2-phenyl-4-cyclohexyl-5-(4-methylsulfonylphenyl)oxazole as COX-2 inhibitors. However, the compounds described in this publication are mainly characterized by 4-fluorophenyl and 4-methylsulfonylphenyl at the 4-position and 5-position of oxazole ring, and do not suggest the compounds having specific substituents in combination, as in the present invention.


Not only in COX-2 inhibitors but also in the field of anti-inflammatory drugs, preferable phenyl substituent for 5-membered heterocyclic ring skeleton has been conventionally considered to be monosubstituted phenyl such as 4-methylsulfonylphenyl and 4-methoxyphenyl, and di-substituted phenyl has been barely tried (e.g., UK Patent No. 1206403).

Disclosure of the Invention

The present inventors have intensively studied with the aim of providing a novel compound having antipyretic activity, analgesic activity and anti-inflammatory activity, which is free of side-effects such as disorders in the digestive tract. Surprisingly, they have found that a compound having a secondary substituent such as halogen atom, in particular, fluorine atom, introduced into phenyl such as 4-lower alkylsulfonylphenyl, 4-aminosulfonylphenyl or 4-lower alkylaminosulfonylphenyl, as a substituent for oxazole, has superior selective inhibitory action on COX-2, which resulted in the completion of the present invention.

That is, the present invention relates to heterocyclic aromatic oxazole compounds as shown in the following (1) to (21), pharmaceutically acceptable salts thereof, intermediate compounds for producing such compounds and pharmaceutical compositions comprising such heterocyclic aromatic oxazole compound.


(1) Heterocyclic aromatic oxazole compounds of the formula (I)



wherein

Z is an oxygen atom;


one of R and R₁ is a group of the formula



wherein R₃ is lower alkyl, amino or lower alkylamino, and R₄, R₅, R₆ and R₇ are the same or different and each is hydrogen atom, halogen atom, lower alkyl, lower alkoxy, trifluoromethyl, hydroxy or amino, provided that at least one of R₄, R₅, R₆ and R₇ is not hydrogen atom, and the other is optionally substituted cycloalkyl, optionally substituted heterocyclic group or optionally substituted aryl; and


R₂ is a lower alkyl or a halogenated lower alkyl, and pharmaceutically acceptable salts thereof.

(2) Heterocyclic aromatic oxazole compounds of the above (1), wherein R₁ is a group of the formula




wherein R₃' is lower alkyl or amino, at least one of R₄', R₅', R₆' and R₇' is halogen atom or lower alkyl and the rest is hydrogen atom or halogen atom, and pharmaceutically acceptable salts thereof.

(3) Heterocyclic aromatic oxazole compounds of the above (1), wherein R₁ is a group of the formula



wherein R₃'' is methyl or amino, R₅'' is fluorine atom and R₆'' is hydrogen atom or fluorine atom, and R₂ is methyl, and pharmaceutically acceptable salts thereof.

(4) Heterocyclic aromatic oxazole compounds of the above (1), wherein R₁ is a group of the formula



wherein R₃'', R₅'' and R₆'' are as defined in the above (3); R is optionally substituted cycloalkyl having 5 to 7 carbon

atoms, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted morpholino, optionally substituted piperazinyl, optionally substituted piperidyl, optionally substituted phenyl, optionally substituted naphthyl or optionally substituted biphenyl, and R_2 is methyl, and pharmaceutically acceptable salts thereof.

(5) Heterocyclic aromatic oxazole compounds of the above (4), wherein R_3 is amino, and pharmaceutically acceptable salts thereof.

(6) Heterocyclic aromatic oxazole compounds of the above (4), wherein R is optionally substituted cycloalkyl having 5 to 7 carbon atoms, optionally substituted phenyl or optionally substituted thienyl, and pharmaceutically acceptable salts thereof.

(7) Heterocyclic aromatic oxazole compounds of the above (4), wherein R is cyclohexyl or 4-fluorophenyl, and R_1 is 4-aminosulfonyl-3-fluorophenyl, 4-aminosulfonyl-3,5-difluorophenyl, 3-fluoro-4-methylsulfonylphenyl or 3,5-difluoro-4-methylsulfonylphenyl, and pharmaceutically acceptable salts thereof.

(8) Heterocyclic aromatic oxazole compounds of the above (1), which are selected from the group of:

4-cyclohexyl-5-(3-fluoro-4-methylsulfonylphenyl)-2-methyloxazole,

5-(4-aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole,

5-(4-aminosulfonyl-3,5-difluorophenyl)-4-cyclohexyl-2-methyloxazole,


4-cyclohexyl-5-(3,5-difluoro-4-methylsulfonylphenyl)-2-methyloxazole,

and


5-(4-aminosulfonyl-3-fluorophenyl)-4-(4-fluorophenyl)-2-methyloxazole,

and pharmaceutically acceptable salts thereof.

(9) Oxime compounds of the following formula (XI')




wherein R_1'' is



wherein R_4 , R_5 , R_6 and R_7 are as defined in the above (1), and R'' is optionally substituted cycloalkyl or optionally substituted aryl.

(10) Oxime compounds of the above (9) wherein R_1'' is 3-fluorophenyl or 3,5-difluorophenyl, and R'' is cyclohexyl or 4-fluorophenyl.


(11) Ketone compounds of the following formula (IV'')




wherein R_1'' and R'' are respectively as defined in the above (9).

(12) Ketone compounds of the above (11) wherein R_1'' is 3-fluorophenyl or 3,5-difluorophenyl, and R'' is cyclohexyl or 4-fluorophenyl.

(13) Ketomethylene compounds of the following formula (IV''')




wherein R''' is optionally substituted cycloalkyl having 5 to 7 carbon atoms, optionally substituted phenyl or optionally substituted thienyl, and R₁''' is a group of the formula



wherein R₃', R₄', R₅', R₆' and R₇' are as defined in the above (2).

(14) Ketomethylene compounds of the above (13) wherein R''' is cyclohexyl, and R₁''' is 4-aminosulfonyl-3-fluorophenyl, 4-aminosulfonyl-3,5-difluorophenyl, 3-fluoro-4-methylsulfonylphenyl or 3,5-difluoro-4-methylsulfonylphenyl.


(15) Ester compounds of the following formula (V)



wherein R, R₁, R₂ and Z are as defined in the above (1).

(16) Ester compounds of the above (15) wherein R is cycloalkyl and R₂ is lower alkyl.

(17) Amide compounds of the following formula (XVIII')



wherein R₁'' and R'' are respectively as defined in the above (9), and Z and R₂ are as defined in the above (1).

(18) Amide compounds of the above (17) wherein R₁'' is 3-fluorophenyl or 3,5-difluorophenyl, R'' is cyclohexyl or 4-fluorophenyl, and R₂ is lower alkyl.

(19) Pharmaceutical compositions comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of the above (1) or a pharmaceutically acceptable salt thereof.

(20) Cyclooxygenase-2 inhibitors comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of the above (1) or a pharmaceutically acceptable salt thereof as an active ingredient.

(21) Anti-inflammatory agents comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of the above (1) or a pharmaceutically acceptable salt thereof as an active ingredient.

As used herein, lower alkyl means an optionally branched alkyl having 1 to 4 carbon atoms, which is exemplified by methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl, with preference given to methyl.

Lower alkylamino is that wherein amino group is substituted by the above-mentioned lower alkyl, and is exemplified by methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, butylamino, isobutylamino, sec-butylamino and tert-butylamino. Preferred are methylamino and dimethylamino.

Halogen atom means chlorine atom, bromine atom, fluorine atom and the like, with preference given to chlorine atom and fluorine atom. Particularly preferred is fluorine atom.

Lower alkoxy is an optionally branched alkoxy having 1 to 4 carbon atoms, which is exemplified by methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy and tert-butoxy, with preference given to methoxy.

Cycloalkyl means a cycloalkyl having 3 to 8 carbon atoms, which is exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, with preference given to cycloalkyl having 5 to 7 carbon atoms, such as cyclopentyl, cyclohexyl and cycloheptyl. Particularly preferred is cyclohexyl.

Heterocyclic group is a 5- or 6-membered aromatic heterocyclic ring, saturated heterocyclic ring or condensed heterocyclic ring of these heterocyclic rings and benzene ring, all having, besides carbon atom, 1 to 3 hetero atoms selected from nitrogen atom, oxygen atom and sulfur atom as atom(s) constituting the ring. Examples thereof include thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, morpholino, piperazinyl, piperidyl, pyranyl, thiopyranyl, pyridyl, benzothienyl, benzofuranyl, indole, 4,5,6,7-tetrahydroindole, 4,5,6,7-tetrahydrobenzothienyl and 4,5,6,7-tetrahydrobenzofuranyl, with preference given to thienyl, furyl, pyrrolyl, morpholino, piperazinyl and piperidyl, and particular preference given to thienyl.

Aryl is, for example, phenyl, naphthyl or biphenyl. Preferred is phenyl.


Halogenated lower alkyl is that wherein lower alkyl is substituted by the above-mentioned halogen atom, and is exemplified by fluoromethyl, chloromethyl, bromomethyl, iodomethyl, difluoromethyl, dichloromethyl, trifluoromethyl, trichloromethyl, fluoroethyl, chloroethyl, difluoroethyl, dichloroethyl, trifluoroethyl, trichloroethyl, tetrachloroethyl, pentafluoroethyl and fluoropropyl, with preference given to fluoromethyl, chloromethyl, dichloromethyl, difluoromethyl, trichloromethyl and trifluoromethyl.

"Optionally substituted" means that the group may be substituted by 1 to 3 substituents wherein said substituents may be the same or different. The position of the substituents is optional and is not particularly limited. Specific examples include lower alkyl such as methyl, ethyl, propyl, isopropyl, butyl and tert-butyl; hydroxy; lower alkoxy such as methoxy, ethoxy, propoxy and butoxy; halogen atom such as fluorine, chlorine and bromine; nitro; cyano; acyl such as formyl, acetyl and propionyl; acyloxy such as formyloxy, acetyloxy and propionyloxy; mercapto; alkylthio such as methylthio, ethylthio, propylthio, butylthio and isobutylthio; amino; alkylamino such as methylamino, ethylamino, propylamino and butylamino; dialkylamino such as dimethylamino, diethylamino, dipropylamino and dibutylamino; carbonyl; alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl; amide; trifluoromethyl; alkylsulfonyl such as methylsulfonyl and ethanesulfonyl; aminosulfonyl; cycloalkyl such as cyclopentyl and cyclohexyl; phenyl; and acylamide such as acetamide and propionylamide. Preferred are hydroxy, lower alkyl, lower alkoxy, mercapto, lower alkylthio, halogen atom, trifluoromethyl, alkylcarbonyl, alkoxycarbonyl and acylamide.

More specifically, optionally substituted aryl means an aryl which may be substituted by halogen atom, hydroxy, lower alkyl, lower alkoxy, lower alkylsulfonyl and aminosulfonyl, particularly phenyl, and is exemplified by phenyl, fluorophenyl, methylphenyl, methoxyphenyl, methylsulfonylphenyl and aminosulfonylphenyl, with preference given to phenyl and 4-fluorophenyl.

Optionally substituted heterocyclic group means a heterocyclic group which may be substituted by halogen atom, hydroxy, lower alkyl, lower alkoxy, lower alkylsulfonyl and aminosulfonyl, and particularly means thienyl, furyl, 5-methylthienyl and 5-chlorothieryl. Optionally substituted cycloalkyl means a cycloalkyl which may be substituted by the same substituents as above, with preference given to cyclohexyl.

Examples of preferable R of the heterocyclic aromatic oxazole compounds of the present invention include cyclohexyl, 4-fluorophenyl and 5-chlorothieryl, with particular preference given to cyclohexyl. Preferred as R₁ is a group of the formula



wherein R₃, R₄, R₅, R₆ and R₇ are as defined above, with particular preference given to a group wherein R₃ is amino

or methyl, R_4 and R_7 are hydrogen atoms and at least one of R_5 and R_6 is fluorine atom. Specific examples include 4-aminosulfonyl-3-fluorophenyl, 3-fluoro-4-methylsulfonylphenyl, 4-aminosulfonyl-3,5-difluorophenyl and 3,5-difluoro-4-methylsulfonylphenyl, with particular preference given to 4-aminosulfonyl-3-fluorophenyl. Preferred as R_2 is methyl.


Pharmaceutically acceptable salt may be any as long as it forms a non-toxic salt with the oxazole derivative of the formula (I). Alkali metal salts such as sodium salt and potassium salt, alkaline earth metal salts such as magnesium salt and calcium salt, ammonium salt, organic base salts such as trimethylamine salt, triethylamine salt, pyridine salt, picoline salt, dicyclohexylamine salt and N,N' -dibenzylethylenediamine salt, and amino acid salts such as lysine salt and arginine salt are among the examples. It may be a hydrate as the case demands.

The compound of the present invention has particularly superior selective inhibitory action on COX-2 and is expected to make a therapeutic drug useful for antipyresis, pain relief and anti-inflammation, which is free of side-effects such as digestive tract disorders.

When the compound of the formula (I) of the present invention or a pharmaceutically acceptable salt thereof is used as a pharmaceutical preparation, it is generally admixed with pharmacologically acceptable carriers, excipients, diluents, extenders, disintegrators, stabilizers, preservatives, buffers, emulsifying agents, aromatics, colorings, sweeteners, thickeners, flavorings, solubilizers and other additives known *per se*, such as water, vegetable oil, alcohol such as ethanol and benzyl alcohol, polyethylene glycol, glycerol triacetate gelatin, carbohydrates such as lactose and starch, magnesium stearate, talc, lanolin and petrolatum, and formulated into, by a conventional method, tablets, pills, powders, granules, suppositories, injections, eye drops, liquids, capsules, troches, aerosols, elixirs, suspensions, emulsions, syrups and the like, which can be administered orally or parenterally.

While the dose varies depending on the kind and severity of the disease, compound to be administered, administration route, and age, sex, body weight etc. of patients, 0.1 mg - 1,000 mg, particularly 1 mg - 300 mg of compound (I) is generally administered orally to an adult per day.

The compounds of the present invention can be produced, for example, by the following methods. It is needless to say that the method for producing the compounds of the present invention is not limited to these methods.



wherein R_2' is lower alkyl or halogenated lower alkyl wherein R_2' may be the same with or different from R_2 , X and X' are the same or different and each is halogen atom such as bromine atom and chlorine atom, X_1 is halogen atom or hydroxy, X_1' is halogen atom or hydroxy or alkali metal derivative thereof, and R , R_1 , R_2 and Z are as defined above.

Step 1

Compound (IV) can be synthesized by reacting compound (II) with compound (III) in the presence of a metal such as zinc and magnesium in an inert solvent such as 1,2-dimethoxyethane, dioxane, ether, tetrahydrofuran, methylene chloride, benzene and toluene at room temperature. In this case, a catalyst such as palladium(O) complex and copper(I) complex may be added.

Step 2

Compound (V) can be synthesized by reacting compound (IV) in acetic acid solvent in the presence of lead tetraacetate, or by refluxing compound (IV) under heating in the presence of a complex such as manganese acetate, in lower alkanecarboxylic acid such as acetic acid and propionic acid corresponding to R_2COOH wherein R_2 is as defined above and benzoic acid and a solvent such as benzene as necessary.

Step 3

Compound (I) can be synthesized by refluxing compound (V) under heating in the presence of ammonium salt (e.g., lower alkanecarboxylic acid ammonium such as ammonium acetate and ammonium formate), and inorganic ammonium such as ammonium carbonate in an acidic solvent such as lower alkanecarboxylic acid (e.g., formic acid, acetic acid and propionic acid). In this reaction, when R or R_1 is aromatic heterocycle, isomers may be produced wherein the 4-position R and the 5-position R_1 are reversed.

Compound (I) can be also synthesized by the following route.

Step 4 wherein X_1 is hydroxy

This step, Step 6 and Step 7 are advantageous when R_2 (e.g., methyl) is converted to other R_2 (e.g., R_2' such as ethyl).

When X_1 is hydroxy, compound (VI) can be synthesized by reacting compound (V) in the presence of a base such as potassium carbonate, lithium hydroxide, sodium hydroxide and potassium hydroxide in an organic solvent such as methanol, ethanol and dioxane, water or a mixed solvent thereof from under cooling to under heating.

Compound (VI) can be also synthesized by the following Step 5.

Step 5 wherein X_1 is halogen atom or hydroxy

Compound (VI) can be synthesized by reacting compound (IV) in the presence of a halogenating agent such as bromine, chlorine and N-bromosuccinimide in an inert solvent such as acetic acid, 1,2-dimethoxyethane, dioxane, ether, tetrahydrofuran, methylene chloride, benzene and toluene to give compound (VI) wherein X_1 is halogen atom. Compound (VI) wherein X_1 is hydroxy can be synthesized by oxidizing compound (IV) with an oxidizing agent such as benzene iodoacetate, or by treating the halogenated compound (VI) obtained above with water in an inert solvent such as acetone, 1,2-dimethoxyethane, dioxane, ether, tetrahydrofuran, benzene and toluene.

Step 6


Compound (V') can be obtained by reacting compound (VI) and compound (VII') by a known method. Specifically, compound (VI) wherein X_1 is hydroxy and compound (VII') wherein X_1' is halogen atom, or compound (VI) wherein X_1 is halogen atom and compound (VII') wherein X_1' is hydroxy are reacted in pyridine, or in the presence of a base such as triethylamine and sodium hydroxide, in an organic solvent such as methylene chloride, chloroform and ethanol, from under cooling to under heating. When X_1 is halogen atom, alkali metal salt such as sodium acetate may be used instead of carboxylic acid compound (VII'). In this case, a base may or may not be added.

Step 7


Compound (I') can be obtained by treating compound (V') in the same manner as in Step 3.

When a compound wherein either R or R_1 is 4-aminosulfonyl-3-fluorophenyl is desired, the compound can be produced from a compound having 3-fluoro-4-methylsulfonylphenyl corresponding to the objective compound by a known method.

Instead of obtaining compound (IV) using, as mentioned above, compound (II) or (III) having, as R or R₁,




wherein R₃, R₄, R₅, R₆ and R₇ are as defined above, compound (II') or (III') having




wherein R₄, R₅, R₆ and R₇ are as defined above, may be used as a starting material to give compound (IV') according to Step 10, which compound is then converted to aminosulfonyl or methylsulfonyl according to the method of Step 15 to give compound (IV). Alternatively, such starting materials (II') and (III') may be used to give a non-sulfonylated oxazole compound (XIII) corresponding to the ultimate compound (I) or (I') according to Step 1 to Step 7, and the obtained compound (XIII) may be subjected to sulfonylation in the same manner as in Step 15 to give the objective compound (I) or (I').

When a compound wherein either R or R₁ is phenyl substituted by alkylaminosulfonyl or aminosulfonyl is desired, compound (X) wherein either R₈ or R₉ is methoxysulfonylphenyl is subjected to the following Step 8 and Step 9 to synthesize compound (IV).



wherein either R₈ or R₉ is methoxysulfonylphenyl of the formula



wherein R₄, R₅, R₆ and R₇ are as defined above, and the other is optionally substituted cycloalkyl, optionally substituted heterocyclic group or optionally substituted aryl, and R, R₁, X and X' are as defined above.

Step 8


Compound (X) can be synthesized in the same manner as in Step 1, using compound (VIII) and compound (IX).

Step 9


When at least one of R and R₁ is phenyl having aminosulfonyl or alkylsulfonyl at the 4-position, compound (IV) can be synthesized by heating compound (X) in pyridine, or refluxing compound (X) under heating in the presence of sodium iodide, potassium iodide, lithium iodide and the like, in an organic solvent such as acetone and tetrahydrofuran, after which the obtained compound is reacted with thionyl chloride or oxalyl chloride under heating. Then, the resulting product is aminated or alkylaminated or alkylated by a known method. More specifically, amination or alkylamination is carried out by reacting the resulting product in the presence of aqueous ammonia or alkylamine, or a base such as sodium acetate and ammonium salt such as alkylamine hydrochloride, in an organic solvent such as tetrahydrofuran, ether, toluene, benzene, methylene chloride and dioxane from under cooling to under heating. The alkylation can be carried out by the method described in J. Org. Chem., 56: 4974-4976 (1991).

Compound (I) can be also synthesized by the method of the following Step 10 to Step 15.

This method is directed to finally introducing sulfonyl group in the last Step 15.



wherein either R' or R₁' is phenyl of the formula



wherein R₄, R₅, R₆ and R₇ are as defined above, and the other is a group corresponding to one of R and R₁, cycloalkyl which may be substituted by a substituent such as lower alkyl, heterocyclic group such as thienyl and furyl, which may be substituted by a substituent lower alkyl or halogen atom, or aryl which may be substituted by a substituent such as halogen atom, lower alkyl and lower alkoxy, and R, R₁, X, X' and Z are as defined above.

Step 10

Compound (IV') can be synthesized in the same manner as in Step 1, wherein compound (II') and compound (III') are reacted in the presence of a metal such as zinc and magnesium in an inert solvent such as 1,2-dimethoxyethane,

dioxane, ether, tetrahydrofuran, methylene chloride, benzene and toluene at room temperature. In this case, a catalyst such as palladium(O) complex and copper(I) iodide complex may be added.

Step 11

Compound (XI) can be synthesized by refluxing under heating compound (IV') and hydroxylamine hydrochloride in the presence of a base such as sodium acetate, sodium hydroxide and potassium carbonate in an organic solvent such as methanol, ethanol and tetrahydrofuran, water or a mixed solvent thereof.

Step 12

Compound (XII) can be synthesized by reacting compound (XI) in the presence of an acylating agent such as acetic anhydride and acetyl chloride, in pyridine, or in the presence of a base such as triethylamine in an organic solvent such as methylene chloride and chloroform from under cooling to under heating.

Step 13

Compound (XIII) can be synthesized by refluxing under heating compound (XII) in an acidic solvent such as formic acid and acetic acid. In this case, a dehydrating agent such as magnesium sulfate and sodium sulfate may be added.

Step 14


This step is for the synthesis of compound (XIII) from compound (XI) in a single step, and compound (XIII) can be synthesized from compound (XI) and carboxylic acid chloride such as acetyl chloride by the method described in Indian J. Chem., 20B: 322-323 (1981). When R₂ is methyl, compound (XIII) can be synthesized by reacting compound (XI) and acetic anhydride while heating in acetic acid.

Step 15

Compound (I) can be synthesized by reacting compound (XIII) in the presence of a chlorosulfonylating agent such as chlorosulfonic acid in an organic solvent such as chloroform and methylene chloride, or without solvent, and subjecting the resulting product to amination, alkylamination or alkylation by a known method. The amination and alkylamination in Step 15 specifically comprise reacting in the presence of aqueous ammonia, alkylamine or a base such as sodium acetate and ammonium salt such as alkylamine hydrochloride in an organic solvent such as tetrahydrofuran, ether, toluene, benzene, methylene chloride and dioxane from under cooling to under heating. When alkylsulfonation is carried out, the method described in J. Org. Chem., 56: 4974-4976 (1991) can be used for the synthesis.

In the above description, alkylsulfonation or aminosulfonation in the final Step 15 has been exemplarily discussed. It is possible to use compound (II) and compound (III) instead of the starting materials (II') and (III') to give compound (IV), which is followed by Step 11 to Step 14 to give an oxazole compound (I). In this case, Step 15 is not necessary.

Compound (XIII) used in Step 15 can be also synthesized by the following route.



wherein R', R₁', R₂ and Z are as defined above.


Step 16

Compound (V'') can be synthesized in the same manner as in Step 2 wherein compound (IV') is reacted in the presence of lead tetraacetate in acetic acid solvent, or by heating compound (IV') in the presence of a complex such as manganese acetate in lower alkanecarboxylic acid such as acetic acid and propionic acid corresponding to R₂COOH wherein R₂ is as defined above, and benzoic acid and in a solvent such as benzene as necessary.

Step 17

Compound (XIII) can be synthesized in the same manner as in Step 3 wherein compound (V'') is refluxed under heating in the presence of ammonium salt such as lower alkanecarboxylic acid ammonium (e.g., ammonium acetate and ammonium formate) and inorganic ammonium (e.g., ammonium carbonate) in an acidic solvent of lower alkanecarboxylic acid such as formic acid, acetic acid and propionic acid. In this reaction, when R' or R₁' is an aromatic heterocycle, isomers may be produced wherein the 4-position R' and the 5-position R₁' are reversed.

Compound (I) can be also synthesized by the method shown in the following Step 18 to Step 21.



wherein X₂ is halogen atom, and R, R₁, R', R₁', R₂ and Z are as defined above.

Step 18

Compound (XV) can be synthesized by reacting compound (XIV) with chlorocarbonate such as ethyl chlorocarbonate in an inert solvent such as tetrahydrofuran, toluene and ethyl acetate in the presence of a base such as triethylamine, or by heating compound (XIV) in acetic anhydride.

Step 19

Compound (XVII) can be synthesized by reacting compound (XV) with compound (XVI) or an acid anhydride corresponding to compound (XVI) in an inert solvent such as tetrahydrofuran, acetonitrile, ethyl acetate and toluene in the presence of magnesium salt such as magnesium chloride and a base such as triethylamine, pyridine and potassium carbonate. Compound (XVII) can be also synthesized by the method described in Chem. Ber., 102: 883-898 (1969).

Step 20

Compound (XVIII) can be synthesized by treating compound (XVII) with an acid such as 1N-4N hydrochloric acid, oxalic acid and dilute sulfuric acid in an inert solvent such as tetrahydrofuran, dioxane, methylene chloride and toluene, or heating compound (XVII) in the presence of pyridine and acetic acid.

Step 21

Compound (I) is obtained by reacting compound (XVIII) with a chlorosulfonylating agent such as chlorosulfonic acid in an organic solvent such as chloroform and methylene chloride, or without solvent. Then, the obtained product is reacted with aqueous ammonia or alkylamine in an organic solvent such as tetrahydrofuran, ether, toluene, methylene

chloride and dioxane, or reacted with ammonium salt such as alkylamine hydrochloride in the presence of a base such as sodium acetate, pyridine and sodium hydroxide.

Compound (I) can be also synthesized from compound (XVIII) by the following Step 22 and Step 23.

Step 22

Compound (XIII) can be synthesized by reacting compound (XVIII) with inorganic acid such as concentrated sulfuric acid and polyphosphoric acid in acetic anhydride, or without solvent, at room temperature to under heating.

Step 23

Compound (I) can be synthesized by reacting compound (XIII) in the same manner as in the aforementioned Step 15.


In the above Step 22 and Step 23, alkylsulfonylation or aminosulfonylation in the final Step 23 has been exemplarily discussed. It is possible to subject a compound having R and R₁ instead of R' and R₁' to the reaction according to Step 18 to Step 20, followed by Step 22 to give an oxazole compound (I). In this case, Step 23 is not necessary.

The compound (I) thus obtained can be isolated and purified by a known method for separation and purification, such as concentration, concentration under reduced pressure, solvent extraction, crystal precipitation, recrystallization and chromatography.

The present invention is described in more detail in the following by illustrative Examples and Experimental Examples, to which the present invention is not limited.


Example 1

Synthesis of 5-(2-chloro-4-methylsulfonylphenyl)-4-cyclohexyl-2-methyloxazole (formula (I'); R=cyclohexyl, R₁=2-chloro-4-methylsulfonylphenyl, R₂'=methyl, Z=oxygen atom) Step 1) 2-Chloro-4-methylsulfonylbenzyl cyclohexyl ketone (formula (IV); R=cyclohexyl, R₁=2-chloro-4-methylsulfonylphenyl)




To a solution of tetrakis(triphenylphosphine)palladium (1.29 g) and zinc powder (2.19 g) in 1,2-dimethoxyethane (10 ml) was added a solution of cyclohexanecarbonyl chloride (3.60 g) in 1,2-dimethoxyethane (10 ml) at room temperature under a nitrogen atmosphere. A solution of 2-chloro-4-methylsulfonylbenzyl bromide (9.40 g) in 1,2-dimethoxyethane (20 ml) was gradually added dropwise to the mixture at room temperature with stirring. The mixture was further stirred at room temperature for 3 hours. The insoluble matter was removed by filtration and the filtrate was concentrated under reduced pressure. Then, ethyl acetate (200 ml) was added to the residue, and the mixture was washed with 1N hydrochloric acid, and then with saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated, and ethyl acetate and diisopropyl ether were added. The precipitated solid was collected by filtration to give 3.47 g of the title compound as a white solid.

Step 5) 2-Bromo-2-(2-chloro-4-methylsulfonylphenyl)-1-cyclohexyl-1-ethanone (formula (VI); R=cyclohexyl, R₁=2-chloro-4-methylsulfonylphenyl, X₁=bromine atom)




To a solution of the compound (3.40 g) obtained in the above Step 1) in benzene (20 ml) was dropwise added a solution of bromine (1.73 g) in benzene (20 ml) with stirring under ice-cooling, and the mixture was stirred for one hour. This solution was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 4.20 g of the title compound.

Step 6) 1-(2-Chloro-4-methylsulfonylphenyl)-2-cyclohexyl-2-oxoethyl acetate (formula (V')); R=cyclohexyl, R₁=2-chloro-4-methylsulfonylphenyl, R₂'=methyl, Z=oxygen atom)



Sodium acetate (1.06 g) and ethanol (40 ml) were added to the compound (4.20 g) obtained in the above Step 5). The mixture was refluxed under heating for 4 hours, and the solvent was evaporated under reduced pressure. Ethyl acetate was added to the residue. The mixture was washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give 3.85 g of a crude product of the title compound.

Step 7) 5-(2-Chloro-4-methylsulfonylphenyl)-4-cyclohexyl-2-methyloxazole (formula (I')); R=cyclohexyl, R₁=2-chloro-4-methylsulfonylphenyl, R₂'=methyl, Z=oxygen atom)




A solution of the compound (3.85 g) obtained in the above Step 6) and ammonium acetate (2.08 g) in acetic acid (40 ml) was refluxed under heating for 5 hours. The solvent was evaporated under reduced pressure, and ethyl acetate was added to the residue. The mixture was washed with water, saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 1.95 g of the title compound (yield 53%).

Example 2


Synthesis of 5-(4-aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole (formula (I)); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂'=methyl, Z=oxygen atom)

Step 10) Cyclohexyl 3-fluorobenzyl ketone (formula (IV')); R'=cyclohexyl, R₁'=3-fluorophenyl)




To a solution of tetrakis(triphenylphosphine)palladium (2.00 g) and zinc powder (17.98 g) in 1,2-dimethoxyethane (50 ml) was added a solution of cyclohexanecarbonyl chloride (20.00 g) in 1,2-dimethoxyethane (50 ml) at room temperature under a nitrogen atmosphere. A solution of 3-fluorobenzyl bromide (26.00 g) in 1,2-dimethoxyethane (100 ml) was gradually added dropwise to the mixture with stirring under ice-cooling. The mixture was stirred under ice-cooling for 30 minutes, and at room temperature for 2 hours. The insoluble matter was removed by filtration and the filtrate was concentrated under reduced pressure. Then, ethyl acetate (200 ml) was added to the residue, and the mixture was washed with 1N hydrochloric acid, and then with saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give 29.20 g of an oily crude product.

Step 16) 2-Cyclohexyl-1-(3-fluorophenyl)-2-oxoethyl acetate (formula (V'));
 R_1 =cyclohexyl, R_1' =3-fluorophenyl, R_2 =methyl, Z=oxygen atom)




Lead tetraacetate (75.00 g) was added to a solution of the compound (29.20 g) obtained in the above Step 10) in acetic acid (300 ml). The mixture was refluxed under heating for 1.5 hours, and the solvent was evaporated under reduced pressure. Ethyl acetate was added to the residue. The mixture was washed with water, a saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue was purified by silica gel column chromatography (developing solvent; hexane:ethyl acetate=9:1) to give 18.30 g of the title compound as an oil (yield 50%).

Step 17) 4-Cyclohexyl-5-(3-fluorophenyl)-2-methyloxazole (formula (XIII));
 R =cyclohexyl, R_1 =3-fluorophenyl, R_2 =methyl, Z=oxygen atom)



A solution of the compound (18.00 g) obtained in the above Step 16) and ammonium acetate (15.00 g) in acetic acid (100 ml) was refluxed under heating for 5 hours, and the solvent was evaporated under reduced pressure. Ethyl acetate was added to the residue. The mixture was washed with water, saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 17.20 g of an oily crude product. Step 15) 5-(4-Aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole (formula (I));
 R =cyclohexyl, R_1 =4-aminosulfonyl-3-fluorophenyl, R_2 =methyl, Z=oxygen atom)



15 To a solution of the compound (17.00 g) obtained in the above Step 17) in chloroform (80 ml) was added dropwise chlorosulfonic acid (27 ml) with stirring under ice-cooling, and the mixture was heated at 100°C for 3 hours. The reaction mixture was cooled to room temperature, and dropwise added to ice-water (300 ml) with stirring. The organic layer was separated, washed with saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 20.31 g of a crude product.

20 Aqueous ammonia (28%) was added to a solution of the obtained compound (10.00 g) in tetrahydrofuran (40 ml) with stirring at room temperature, and the mixture was stirred at room temperature for one hour. The solvent was evaporated under reduced pressure and ethyl acetate was added to the residue. The mixture was washed with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated, and the residue was separated and purified by silica gel column chromatography (developing solvent; dichloromethane:ethyl acetate=6:1) to give 5.74 g of the title compound (yield 61%).

25 Example 2'


The compound of Example 2 (formula (I); R=cyclohexyl, R_1 =4-aminosulfonyl-3-fluorophenyl, R_2 =methyl, Z=oxygen atom) was synthesized according to another synthetic method.

30 Step 11) Cyclohexyl 3-fluorobenzyl ketone oxime (formula (XI); R' = cyclohexyl, R_1' =3-fluorophenyl)



45 To a solution of the compound (353 g) obtained according to a method similar to that of the above Example 2, Step 10) in ethanol (1300 ml) were added hydroxylamine hydrochloride (123 g) and sodium acetate (158 g). The mixture was refluxed under heating for 2 hours, and the solvent was evaporated under reduced pressure. Ethyl acetate was added to the residue. The mixture was washed with water, saturated aqueous sodium hydrogencarbonate solution and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the crude product was recrystallized from n-heptane to give 160 g of the title compound (yield 42%).

Step 14) 4-Cyclohexyl-5-(3-fluorophenyl)-2-methyloxazole (formula (XIII); R' =cyclohexyl, R_1' =3-fluorophenyl, R_2 =methyl, Z=oxygen atom)




Acetic anhydride (95 ml) was dropwise added to a solution of the compound (158 g) obtained in the above Step 11) in acetic acid (900 ml) with stirring at room temperature, and the mixture was refluxed under heating for 7 hours. The solvent was evaporated under reduced pressure and n-heptane was added to the residue. The mixture was washed with water, saturated aqueous sodium hydrogencarbonate solution, saturated brine and acetonitrile. The solvent was evaporated under reduced pressure to give 119 g of the title compound as an oil.

Then, the obtained compound (119 g) was reacted in the same manner as in the above Example 2, Step 15) to give a compound of Example 2 (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom).

Example 3

Synthesis of 4-cyclohexyl-5-(3-fluoro-4-methylsulfonylphenyl)-2-methyloxazole (formula (I); R=cyclohexyl, R₁=3-fluoro-4-methylsulfonylphenyl, R₂=methyl, Z=oxygen atom)

Step 15) 4-Cyclohexyl-5-(3-fluoro-4-methylsulfonylphenyl)-2-methyloxazole (formula (I); R=cyclohexyl, R₁=3-fluoro-4-methylsulfonylphenyl, R₂=methyl, Z=oxygen atom)



To a solution of the compound (17.00 g) obtained in the above Example 2, Step 17) in chloroform (80 ml) was dropwise added chlorosulfonic acid (27 ml) with stirring under ice-cooling. The mixture was heated at 100°C for 3 hours. The reaction mixture was cooled to room temperature and dropwise added to ice-water (300 ml) with stirring. The organic layer was separated, washed with saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 20.31 g of a crude product.

Water (25 ml) was added to the obtained compound (3.66 g). To the mixture were added sodium sulfite (1.42 g) and sodium hydrogencarbonate (1.89 g) successively with stirring at room temperature. The mixture was heated at 70°C for 2 hours. Ethanol (25 ml) and methyl iodide (2.20 g) were added to the mixture, and the mixture was heated at 100°C for 2 hours. The mixture was cooled to room temperature and extracted with ethyl acetate. The extract was washed with saturated brine and dried over anhydrous sodium sulfate.

The solvent was evaporated under reduced pressure, and the residue was separated and purified by silica gel column chromatography (developing solvent; hexane:ethyl acetate=2:1) to give 0.82 g of the title compound (yield 24%).

Examples 4-6

The compounds of Examples 4-6 were obtained in the same manner as in Examples 1-3 or Example 7 to be mentioned below.

The structures and properties of the compounds of Examples 1-6 are shown in the following Tables. In the Tables, Me means methyl.

Table 1








Ex.	Compound	m.p.	¹ H NMR (δ) ppm	IR cm ⁻¹	MS	Elem. analysis
1		119 ~ 121 °C white crystals	CDCl ₃ 300MHz 1.1 - 1.2 (3H, m) 1.6 - 1.8 (7H, m) 2.48 (1H, m) 2.51 (3H, s) 3.12 (3H, s) 7.55 (1H, d, J=8.1Hz) 7.88 (1H, dd, J=1.8, 8.1Hz) 8.07 (1H, d, J=1.8Hz)	neat 2928 1578 1317 1155 1100 960	FAB+ 354 (MH ⁺)	
2		166 ~ 167 °C white crystals	CDCl ₃ 300MHz 1.3 - 1.5 (3H, m) 1.6 - 1.9 (7H, m) 2.51 (3H, s) 2.79 (1H, t, J=3.7, 11.3Hz) 5.11 (2H, s) 7.36 - 7.44 (2H, m) 7.94 (1H, t, J=7.9Hz)	neat 3280 2929 1613 1343 1170	FAB+ 339 (MH ⁺)	Calculated C 56.79 % H 5.66 % N 8.28 % Found C 56.41 % H 5.73 % N 8.19 %
3		111 ~ 112 °C white crystals	CDCl ₃ 300MHz 1.3 - 1.5 (3H, m) 1.6 - 1.8 (7H, m) 2.52 (3H, s) 2.80 (1H, t, J=4.0, 11.4Hz) 3.25 (3H, s) 7.40 (1H, dd, J=1.6, 11.2Hz) 7.48 (1H, dd, J=1.6, 8.3Hz) 7.99 (1H, dd, J=8.3, 8.4Hz)	neat 2929 1612 1320 1161 1144 769	FAB+ 338 (MH ⁺)	Calculated C 60.52 % H 5.97 % N 4.15 % Found C 60.70 % H 6.10 % N 4.12 %

Table 2

Ex.	Compound	m.p.	¹ H NMR (δ) ppm	IR cm ⁻¹	MS	Elem. analysis
4		200 ~ 201 °C white crystals	CDCl ₃ 300MHz 1.28 - 1.44 (4H, m) 1.62 - 1.92 (6H, m) 2.51 (3H, s) 2.72 - 2.83 (1H, m) 5.18 (2H, s) 7.53 (1H, dd, J=8.4, 1.6Hz) 7.69 (1H, d, J=1.6Hz) 8.13 (1H, d, J=8.4Hz)	KBr 3353 3255 2928 1606 1342 1166	FAB+ 355 (MH ⁺)	Calculated C 54.16 % H 5.40 % N 7.89 % Found C 54.11 % H 5.45 % N 7.78 %
5		183.2 ~ 184.2 °C white crystals	CDCl ₃ 300MHz 1.3 - 1.5 (3H, m) 1.7 - 1.9 (7H, m) 2.50 (3H, s) 2.73 (3H, s) 2.80 (1H, m) 4.92 (2H, s) 7.43 - 7.49 (2H, m) 8.05 (1H, d, J=8.3Hz)	KBr 3294 2929 1609 1299 1170	FAB+ 335 (MH ⁺)	Calculated C 61.05 % H 6.63 % N 8.38 % Found C 61.24 % H 6.73 % N 8.43 %
6		amorphous	CDCl ₃ 300MHz 1.28 - 1.47 (3H, m) 1.57 - 1.95 (7H, m) 2.51 (3H, s) 2.68 - 2.80 (1H, m) 5.37 (2H, brs) 7.18 (2H, ddd, J=9.9, 1.7, 1.4Hz)	KBr 2931 1622 1557 1422 1359 1175 1035	FAB+ 357 (MH ⁺)	

Example 7


Synthesis of 5-(4-aminosulfonyl-3-fluorophenyl)-4-(4-fluorophenyl)-2-methyloxazole (formula (I)); R=4-fluorophenyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom)



A solution of 5-(3-fluorophenyl)-4-(4-fluorophenyl)-2-methyloxazole (1.10 g) obtained by the method as mentioned above and chlorosulfonic acid (1.6 ml) in chloroform (2 ml) was heated with stirring at 90°C for 2 hours. The reaction mixture was poured into ice-water and extracted with chloroform. The organic layer was washed with water and saturated brine, dried over anhydrous magnesium sulfate, and concentrated to give 1.06 g of a crude product of 5-(4-chlorosulfonyl-3-fluorophenyl)-4-(4-fluorophenyl)-2-methyloxazole.

To a solution of this crude product (1.06 g) in tetrahydrofuran (6 ml) was added 28% aqueous ammonia (0.6 ml) and the mixture was stirred at room temperature for 2 hours. The reaction mixture was concentrated, added with ethyl acetate, and washed with water and saturated brine. The ethyl acetate solution was dried over anhydrous magnesium sulfate, and concentrated to give 981 mg of a crude product. This crude product was recrystallized from ethanol to give 629 mg of the title compound (yield 44%). The structure and properties of this compound are shown in the following Table.


Table 3

Ex.	Compound	m. p.	¹ H NMR (δ) ppm	IR cm ⁻¹	MS	Elem. analysis
7		208 °C white crystals	CDCl ₃ , 300MHz 2.58 (3H, s) 5.07 (2H, s) 7.14 (2H, t, J=2.2, 8.8Hz) 7.36 (1H, dd, J=1.5, 11.0Hz) 7.47 (1H, dd, J=1.8, 7.7Hz) 7.59 (2H, ddd, J=2.2, 5.5, 8.8Hz) 7.88 (1H, t, J=7.7Hz)	neat 3278 2359 1613 1562 1510 1342 1171	FAB+ 351 (M ⁺ +1)	Calculated C 54.74 % H 3.86 % N 7.66 % Found C 54.40 % H 3.74 % N 7.59 %

Example 2"


The compound of Example 2 (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom) was synthesized according to another synthetic method.

Step 18) 4-Cyclohexyl-2-methyl-5-oxazolone (formula (XV); R'=cyclohexyl, R₂=methyl)




Triethylamine (8.39 ml) was added to a suspension of DL-N-acetyl-2-cyclohexylglycine (10.00 g) obtained from α -aminophenylacetic acid according to a known method [Collect. Czech. Chem. Commun., 31: 4563 (1996)] in ethyl acetate (50 ml). Ethyl chlorocarbonate (5.28 ml) was dropwise added to the mixture under ice-cooling. The mixture was stirred under ice-cooling for one hour, added with ethyl acetate (150 ml), and washed successively with water and saturated brine. The ethyl acetate solution was concentrated under reduced pressure to give 9.86 g of the title compound as an oil.

Step 19) 4-Cyclohexyl-4-(3-fluorobenzoyl)-2-methyl-5-oxazolone (formula (XVII); R'=cyclohexyl, R₁'=3-fluorophenyl, R₂=methyl, Z= oxygen atom)




A solution of the compound (9.86 g) obtained in the above Step 18) in tetrahydrofuran (15 ml) was added to a suspension of magnesium chloride (3.56 g) in tetrahydrofuran (20 ml). Triethylamine (9.49 ml) was added with stirring under ice-cooling, and the mixture was stirred for 15 minutes. 3-Fluorobenzoyl chloride (4.55 ml) was dropwise added to the mixture, and the mixture was stirred under ice-cooling for one hour. The reaction mixture was diluted with ethyl acetate, washed with water, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 11.69 g of the title compound as an oil.

Step 20) 2-N-Acetylamino-2-cyclohexyl-3'-fluoroacetophenone (formula (XVIII); R' =cyclohexyl, R₁'=3-fluorophenyl, R₂=methyl, Z=oxygen atom)



To a solution of the compound (527 mg) obtained in the above Step 19) in tetrahydrofuran (3.5 ml) was added 1N hydrochloric acid (0.35 ml). The mixture was stirred at room temperature for one hour, added with ethyl acetate, and washed successively with water, saturated aqueous sodium hydrogencarbonate solution and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give 404 mg of the title compound as a solid (yield 84%). The solid was recrystallized from n-heptane to give white crystals, melting point 116-117°C. Step 21) 5-(4-Aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom)




Chlorosulfonic acid (0.34 ml) was added to a solution of the compound (200 mg) obtained in the above Step 20) in chloroform (2 ml) with stirring under ice-cooling, and the mixture was refluxed under heating for 5 hours. The reaction mixture was diluted with chloroform and poured into ice-water. The organic layer was separated, washed successively with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 181 mg of a crude product.

To a solution of the obtained compound (169 mg) in tetrahydrofuran (2 ml) was added 28% aqueous ammonia (0.1 ml) with stirring at room temperature, and the mixture was stirred for 30 minutes. The solvent was evaporated under reduced pressure. Ethyl acetate was added to the residue, and the mixture was washed successively with water and saturated brine, which was followed by drying over anhydrous sodium sulfate. The solvent was evaporated, and the residue was separated and purified by silica gel column chromatography (developing solvent; dichloromethane:ethyl acetate= 6:1) to give 126 mg of the title compound (yield 55%).

Example 2'''


The compound of Example 2 (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom) was synthesized according to another synthetic method.

Step 22) 4-Cyclohexyl-5-(3-fluorophenyl)-2-methyloxazole (formula (XII); R'=cyclohexyl, R₁'=3-fluorophenyl, R₂=methyl)



Concentrated sulfuric acid (30 µl) was added to a suspension of the compound (141 mg) obtained in the above Example, Step 20) in acetic anhydride (2 ml), and the mixture was stirred at 100°C for 30 minutes. The reaction mixture was concentrated under reduced pressure, added with aqueous potassium carbonate solution, and extracted with ethyl acetate. The organic layer was washed with water and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give 135 mg of the title compound as an oil.

Step 23) 5-(4-Aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom)



15 In the same manner as in the above Example 2, Step 15), the compound obtained in the above Step 22) was reacted to give the compound of Example 2 (formula (I); R=cyclohexyl, R₁=4-aminosulfonyl-3-fluorophenyl, R₂=methyl, Z=oxygen atom).

Experimental Example 1 (inhibitory action on cyclooxygenase)

20 The enzymatic activity was determined from the percent conversion of ¹⁴C arachidonic acid into prostaglandin H₂ (PGH₂) and the decomposed product thereof. That is, a test sample (20 μl), an enzyme solution (20 μl) and distilled water (10 μl) were added to 100 mM Tris-HCl buffer (pH 8, 140 μl) containing hematin (2 μM) and tryptophan (5 mM), and the mixture was thoroughly stirred, which was followed by preincubation at 24°C for 5 minutes. Then, a ¹⁴C arachidonic acid solution (10 μl) was added and the mixture was reacted at 24°C, whereafter a solution (40 μl) of ethyl ether/methanol/1M citric acid (30/4/1) ice-cooled to -20°C was added to stop the reaction. The reaction mixture was centrifuged for 5 minutes at 3,000 rpm to give an ether layer which was placed on a thin plate, and developed with ethyl ether/methanol/acetic acid (90/2/0.1) to determine percent conversion (A) from arachidonic acid to PGH₂ and the decomposed product thereof. The percent conversion (B) without a test sample was also determined, based on which percent inhibition was calculated from the following formula, and a concentration (IC₅₀) necessary for 50% inhibition of the test sample was determined.









$$\text{Inhibition (\%)} = (1 - A/B) \times 100$$

35 An enzyme prepared from human platelets was used as an enzyme solution of cyclooxygenase-1, and an enzyme expressed by a yeast, into which cDNA of human cyclooxygenase-2 had been introduced using a kit of Invitrogen Corp., was used as an enzyme solution of cyclooxygenase-2. As used herein, control compound 1 was 5-(4-aminosulfonylphenyl)-4-cyclohexyl-2-methyloxazole, a patent application to which has been previously filed by us, and control compound 2 was a known analogous compound, 5-(4-aminosulfonylphenyl)-4-(4-fluorophenyl)-2-methyloxazole.

The results are shown in Table 4.

40 As is evident from the comparison of control compound 1 and the compound of Example 2, as well as control compound 2 and the compound of Example 7, a remarkable reduction of the action on COX-1 while retaining the activity on COX-2 has become possible particularly by introducing fluorine atom.

Table 4 : Experimental Example 1 (inhibitory action on cyclooxygenase)

Example	Structural formula	IC ₅₀ (μ M)		COX-1/COX-2
		COX-2	COX-1	
2		0.07	>100	>1,428
3		0.3	>100	>333
4		>10		
5		>10		
6		0.16	>100	>625
7		0.03	37	1,233
Indomethacin		8	0.5	0.063
Control 1		0.07	45	643
Control 2		0.02	5	250

Experimental Example 2 (effects on carrageenin-induced podedema)

Carrageenin (1%, 0.05 ml) dissolved in physiological saline was subcutaneously injected to the left hindlimb of male Donryu rats to induce podedema. The degree of podedema was evaluated by measuring the volume of the limb

3 hours after carrageenin administration. A test compound (1, 3, 10 or 30 mg/kg) was orally administered one hour before carrageenin administration, and suppression thereby was studied. Inhibitory activity was expressed by the dose (ED₃₀) of the test compound necessary for inhibiting by 30% relative to the control group. The results are shown in Table 5.

Table 5

Experimental Example 2 (effects on carrageenin-induced podedema in rats)	
Example	carrageenin-induced podedema in rats, ED ₃₀ (mg/kg p.o.)
2	5.5
indomethacin	2.9


Industrial Applicability

The compound of the present invention, in particular, a compound wherein R₃ is methyl or amino, R₅ is fluorine atom, R₆ is hydrogen atom or fluorine atom, and R₄ and R₇ are hydrogen atom, and pharmaceutically acceptable salts thereof surprisingly selectively inhibit COX-2 alone, while scarcely inhibiting COX-1. Accordingly, the compound of the present invention possesses superior antipyretic action, analgesic action and anti-inflammatory action that the conventional products cannot afford, and scarcely show side-effects in the digestive tract.

Consequently, the development of a superior anti-inflammatory agent heretofor not existed has been enabled, which in turn produces great expectation of the provision of a practical therapeutic agent for the diseases possibly caused by COX-2 product, such as asthma and rheumatism.

Claims


1. A heterocyclic aromatic oxazole compound of the formula (I)



wherein

Z is an oxygen atom;


one of R and R₁ is a group of the formula



wherein R_3 is lower alkyl, amino or lower alkylamino, and R_4 , R_5 , R_6 and R_7 are the same or different and each is hydrogen atom, halogen atom, lower alkyl, lower alkoxy, trifluoromethyl, hydroxy or amino, provided that at least one of R_4 , R_5 , R_6 and R_7 is not hydrogen atom, and the other is optionally substituted cycloalkyl, optionally substituted heterocyclic group or optionally substituted aryl; and


R_2 is a lower alkyl or a halogenated lower alkyl, or a pharmaceutically acceptable salt thereof.

2. The heterocyclic aromatic oxazole compound of Claim 1, wherein R_1 is a group of the formula




wherein R_3' is lower alkyl or amino, at least one of R_4' , R_5' , R_6' and R_7' is halogen atom or lower alkyl and the rest is hydrogen atom or halogen atom, or a pharmaceutically acceptable salt thereof.

3. The heterocyclic aromatic oxazole compound of Claim 1, wherein R_1 is a group of the formula



wherein R_3'' is methyl or amino, R_5'' is fluorine atom and R_6'' is hydrogen atom or fluorine atom, and R_2 is methyl, or a pharmaceutically acceptable salt thereof.

4. The heterocyclic aromatic oxazole compound of Claim 1, wherein R_1 is a group of the formula




wherein R_3'' , R_5'' and R_6'' are as defined in Claim 3; R is optionally substituted cycloalkyl having 5 to 7 carbon atoms, optionally substituted thienyl, optionally substituted furyl, optionally substituted pyrrolyl, optionally substituted morpholino, optionally substituted piperazinyl, optionally substituted piperidyl, optionally substituted phenyl, optionally substituted naphthyl or optionally substituted biphenyl, and R_2 is methyl, or a pharmaceutically acceptable salt thereof.

5. The heterocyclic aromatic oxazole compound of Claim 4, wherein R_3'' is amino, or a pharmaceutically acceptable salt thereof.
6. The heterocyclic aromatic oxazole compound of Claim 4, wherein R is optionally substituted cycloalkyl having 5 to 7 carbon atoms, optionally substituted phenyl or optionally substituted thienyl, or a pharmaceutically acceptable salt thereof.


7. The heterocyclic aromatic oxazole compound of Claim 4, wherein R is cyclohexyl or 4-fluorophenyl, and R₁ is 4-aminosulfonyl-3-fluorophenyl, 4-aminosulfonyl-3,5-difluorophenyl, 3-fluoro-4-methylsulfonylphenyl or 3,5-difluoro-4-methylsulfonylphenyl, or a pharmaceutically acceptable salt thereof.

8. The heterocyclic aromatic oxazole compound of Claim 1, which is selected from the group consisting of:
 4-cyclohexyl-5-(3-fluoro-4-methylsulfonylphenyl)-2-methyloxazole,
 5-(4-aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyloxazole,
 5-(4-aminosulfonyl-3,5-difluorophenyl)-4-cyclohexyl-2-methyloxazole,
 4-cyclohexyl-5-(3,5-difluoro-4-methylsulfonylphenyl)-2-methyloxazole,
 and
 5-(4-aminosulfonyl-3-fluorophenyl)-4-(4-fluorophenyl)-2-methyloxazole,
 or a pharmaceutically acceptable salt thereof.

9. An oxime compound of the following formula (XI')




wherein R₁'' is



wherein R₄, R₅, R₆ and R₇ are as defined in Claim 1, and R'' is optionally substituted cycloalkyl or optionally substituted aryl.

10. The oxime compound of Claim 9 wherein R₁'' is 3-fluorophenyl or 3,5-difluorophenyl, and R'' is cyclohexyl or 4-fluorophenyl.


11. A ketone compound of the following formula (IV'')




wherein R₁'' and R'' are respectively as defined in Claim 9.

12. The ketone compound of Claim 11, wherein R₁'' is 3-fluorophenyl or 3,5-difluorophenyl, and R'' is cyclohexyl or 4-fluorophenyl.

13. A ketomethylene compound of the following formula (IV''')




10 wherein R''' is an optionally substituted cycloalkyl having 5 to 7 carbon atoms, an optionally substituted phenyl or an optionally substituted thienyl, and R1''' is a group of the formula



wherein R3', R4', R5', R6' and R7' are as defined in Claim 2.

14. The ketomethylene compound of Claim 13, wherein R''' is cyclohexyl, and R1''' is 4-aminosulfonyl-3-fluorophenyl, 4-aminosulfonyl-3,5-difluorophenyl, 3-fluoro-4-methylsulfonylphenyl or 3,5-difluoro-4-methylsulfonylphenyl.


15. An ester compound of the following formula (V)



40 wherein R, R1, R2 and Z are as defined in Claim 1.

16. The ester compound of Claim 15, wherein R is cycloalkyl and R2 is lower alkyl.

17. An amide compound of the following formula (XVIII')



55 wherein R1'' and R'' are respectively as defined in Claim 9, and Z and R2 are as defined in Claim 1.

18. The amide compound of Claim 17, wherein R1'' is 3-fluorophenyl or 3,5-difluorophenyl, R'' is cyclohexyl or 4-fluorophenyl, and R2 is lower alkyl.

19. A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of Claim 1 or a pharmaceutically acceptable salt thereof.
20. A cyclooxygenase-2 inhibitor comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of Claim 1 or a pharmaceutically acceptable salt thereof as an active ingredient.
21. An anti-inflammatory agent comprising a pharmaceutically acceptable carrier, and a heterocyclic aromatic oxazole compound of Claim 1 or a pharmaceutically acceptable salt thereof as an active ingredient.

10

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP95/02600

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl. ⁶ C07D263/32, C07D413/04, A61K31/42, C07D207/333, 265/30, 295/10, 307/46, 333/22, C07C251/48, 49/782, 49/792, 49/813, 317/14, 311/16, 311/29, According to International Patent Classification (IPC) or to both national classification and IPC 233/61, 233/76		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int. Cl. ⁶ C07D263/32, C07D413/04, A61K31/42, C07D207/333, 265/30, 295/10, 307/46, 333/22, C07C251/48, 49/782, 49/792, 49/813, 317/14, 311/16, 311/29, 233/61, 233/76 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, 94/27980, A1 (P. D. SEARLE & Co.), December 8, 1994 (08. 12. 94) & US, 5380738, A	1 - 21
X	US, 4782058, A (Pennwalt Corporation), November 1, 1988 (01. 11. 88), Particularly refer to pages 8, 5 (Family: none)	9, 11
X	JP, 62-138485, A (CIBA-Geigy AG.), June 22, 1987 (22. 06. 87), Particularly refer to compound of formula (III) & EP, 225290, A & US, 4849007, A	11
X	JP, 59-155365, A (Shionogi & Co., Ltd.), September 4, 1984 (04. 09. 84), Particularly refer to page 3 & EP, 117578, A & GB, 2136800, A	11
X	J. Org. Chem., Vol. 43, No. 15, (1978) Paul D. Seemuch, et al. "α-Hetero-Substituted Phosphonate Carbanions. 7. Synthesis of Deoxy	11
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search March 6, 1996 (06. 03. 96)		Date of mailing of the international search report April 2, 1996 (02. 04. 96)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP95/02600

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	benzoins and Benzo(b)furans" p. 3063-3065 particularly refer to compound of (3)	
X	J. Org. Chem., Vol. 53, No. 24, (1988) Scott C. Berk, et al. "General approach to highly functionalized benzylic organometallics of zinc and copper" p. 5789-5791 particularly refer to page 5790	11
X	J. Am. Chem. Soc., Vol. 115, No. 8 (1993) Taehee Noh et al. "Photochemistry of α -(-Tolyl) acetone and some derivatives: Triplet α -cleavage and singlet δ -hydrogen abstraction" p. 3105-3110 particularly refer to page 3106	11

Form PCT/ISA/210 (continuation of second sheet) (July 1992)